



ISSN 0375-1511

Rec. zool. Surv. India : 114(Part-3) : 379-390, 2014

FREE LIVING AND PLANT PARASITIC SOIL NEMATODES (ORDERS DORYLAIMIDA AND TYLENCHIDA) OF HIMACHAL PRADESH, INDIA

DEBABRATA SEN, V. V. GANTAIT AND S. BANERJEE

Zoological Survey of India, M- Block, New Alipore, Kolkata - 700053

INTRODUCTION

Himachal Pradesh is a state in Northern part of India with Shimla as the state capital. It is spread over 21,495 sq miles (55,670 km²), and is bordered by the states of Jammu and Kashmir on the north, Punjab on the west and south-west, Haryana and Uttar Pradesh on the south, Uttarakhand on the south-east. Himachal Pradesh is located between 30°10' to 33°12' north latitude and 75°47' to 79°04' east longitude. The state comprises of twelve districts – Shimla, Solan, Chamba, Hamirpur, Bilaspur, Kangra, Kullu, Mandi, Sirmaur, Una, Kinnaur and Lahaul and Spiti. There are major divisions: the middle or central Himalayas comprising of Kinnaur, Kullu, Lahaul and Spiti and Pangi areas, the lesser Himalayas of Dhauladhar ranges and Shimla hills, Shivaliks of Sirmour, Bilaspur, Solan and Hiamirpur districts and river terraces of different valleys like Kullu valley, and Una districts. It is a mountainous state with altitudes ranging from 350 to 69750 meters above the sea level. This wide variation in altitude with mountains, hills and valleys provides different climatic conditions leading to the cultivation of varieties of crops. Based on altitude, rainfall, temperature, humidity and topography, Himachal Pradesh has been divided into four agro-climatic Zones: subtropical sub-mountain and low hills, sub-temperate sub-humid hills, wet temperate high hills and dry temperate high hills and cold desert. The temperature and relative humidity range from 0-35°C and 50-80% respectively. Rainfall varies in between 350-1500 mm in different areas. Soil

type is mostly loamy sand, sandy loam, clay loam and silt loam.

Himachal Pradesh is a land of jade forests and fresh air. As much as 68% of the land area is covered with jungles. While the foothills and valleys are a refreshing green, the areas above the snow line are almost bare and desert type. The southernmost tracts are dominated by sal (*Shorea robusta*), sisham, chir pine, dry deciduous and moist broad-leaved forests. The temperate region above this grows oaks, deodar, blue pine, fir and spruce. In the uppermost climes, trees are sturdy with a vast network of roots (to help them tide over the weeks of burial under heavy snow). Alders, birches, rhododendrons and moist alpine scrubs are found in the name of vegetation. The tough rhododendron is an amazing plant of terrific importance in the ecological chain. About 70-80% population of the state are dependent on farming. The major agricultural crops are sub-tropical and temperate fruits among which apple is one of the most economically important fruit crop. Maize, rice, wheat, ginger, cucurbits, peas, potato, tomato, barley are grown in different seasons.

Among the soil-borne pests, soil free-living and plant-parasitic nematodes are generally overlooked due to their hidden nature and microscopic size. They multiply in millions and spread from place to place by different agricultural practices, causing great damage to the agricultural crops. They inhibit root growth, growth of plants affecting crop production and are thus responsible for massive yield losses. Due to this lack of awareness in

common people and farmers; the importance of nematodes in agriculture was overlooked in this state till 1951. Thirumalachar (1951) first recorded the occurrence of root-knot nematode on potato tubers for the first time from Shimla.

After 1951, extensive studies on taxonomy of tylenchid nematodes were done and several new species from Himachal Pradesh were described by different nematologists (Jairajpuri and Siddiqi, 1963a; Khan, 1964; Jairajpuri & Baqri, 1973; Saha *et al.*, 1973; Khan & Singh, 1974; Jairajpuri and Siddiqi, 1979; Sultan & Jairajpuri, 1978, 1979; Sultan, 1980; Khan & Khan, 1982; Khurma & Gupta, 1988 a, b; Sharma *et al.*, 1986). On the other hand, the study on nematodes belonging to the order Dorylaimida from Himachal Pradesh is comparatively less, although several new species have been described from the state (Jairajpuri & Siddiqi, 1963b; Ali *et al.*, 1974; Ahmad & Jairajpuri, 1979, 1980 and 1982; Baqri & Jairajpuri, 1974, 1975; Jairajpuri & Coomans, 1977).

Nematodes associated with different fruit plants in Himachal Pradesh have been studied by Mukhopadhyaya (1970), Bhardwaj & Sharma (1971), Chandel (1986) and by Sharma & Kaur (1986, 1987). Association of nematodes with mushroom was observed by Bhardwaj *et al.*, (1973) and Chandel (1982). A detailed information on plant nematology of Himachal Pradesh has been provided (Sharma & Gupta, 1998). Extensive work on the host record, occurrence, distribution and association of nematodes with different economically important agricultural and horticultural crops was done (Thirumalachar, 1951; Mukhopadhyaya, 1970; Dalal & Bhatti, 1983; Sharma *et al.*, 1984; Kaur, 1987; Sharma *et al.*, 1988; Khurma, 1989; Kaur *et al.*, 1989; Kaur *et al.*, 1990; Kaur & Sharma, 1990; Chandel, 1993)

The present study includes two systematic lists of soil free-living and plant-parasitic nematodes belonging to the orders Dorylaimida Pearse, 1942 and Tylenchida Thorne, 1949 from Himachal Pradesh. 34 species under 18 genera and 9 families of Dorylaimida and 83 species under 35 genera and

13 families of Tylenchida have been compiled and reported in this report. Among these, *Dorylaimus neominimus* Gantait *et al.* 2010, *Aporcelaimellus heynsi* Baqri and Jairajpuri, 1968., *Labronema glandosum* Rahman *et al.*, 1986, *Thonus garhwaliensis* Ahmad *et al.*, 1986, and the genera *Actinolaimoides* Meyl, 1957, *Laimydorus* Siddiqi, 1969 and *Indokochinema* Darekar and Khan, 1979 are new records from Himachal Pradesh. *Dorylaimus thornei* Andrassy, 1969 is the first report from India. Five species of Tylenchida i.e. *Polenchus shamimi* Baqri, 1991, *Rotylenchoides neoformis* (Siddiqi and Husain, 1964) Sher, 1966, *Rotylenchulus reniformis* Linford and Oleiveira, 1940. *Tylenchorhynchus cylindricus* Cobb, 1913 and the genus *Filenchus* (Meyl, 1961) Andrassy, 1954 are new records from Himachal Pradesh. The respective species of the genera *Actinolaimoides*, *Laimydorus*, *Indokochinema* and *Filenchus* could not be confirmed. The classification proposed by Jairajpuri and Ahmad (1992) and by Siddiqi (2000) have been followed to arrange the available genera and species of Dorylaimida and Tylenchida respectively from Himachal Pradesh and to indicate their present taxonomic status.

MATERIALS AND METHODS

Processing of soil samples and extraction of nematodes:

The collected soil samples were processed by Cobb's sieving and decantation technique (Cobb, 1918) followed by modified Baermann funnel technique (Christie and Perry, 1951) for extraction of nematodes. Soil sample of about 500gms was taken in a bucket of 10 liter capacity. The bucket was filled with clean water up to half of its capacity. The soil and water were thoroughly mixed by hand to make a homogeneous suspension. Stones and plant debris were removed. The muddy suspension was stirred very well by hand and left undisturbed for 20-30 seconds allowing the heavier particles to settle down at the bottom of the bucket while the nematodes and fine soil particles remained suspended in water. This suspension was passed through a coarse sieve and was collected in another bucket. The entire process was repeated thrice to

get a muddy suspension quite free from stones, large soil particles and other undesirable organic matter. This suspension was subjected to similar filtration, but through a fine sieve of 325 meshes. Most of the fine soil particles passed through this sieve but the nematodes and larger soil particles were retained on the surface of the sieve. This was again washed thoroughly with running water within the sieve to get rid of the soil particles as far as possible without losing the nematodes. Then the entire content of the sieve was collected in a beaker of 250 ml, by washing the sieve repeatedly with water, so that all nematodes could be collected.

Then a double layer of tissue paper was placed on an aluminium net of 2 mm pore size and was made moist by applying water gently without leaving any air gap or bubble between the layers of the tissue paper. Now, the aliquot, previously collected in the beaker, was poured gently on this moist double tissue paper supported by 2 mm pore size aluminium net and was further suspended in clean water in a petri dish. The aliquot on the tissue paper was covered by another petri dish to prevent evaporation and desiccation. The whole arrangement was left undisturbed for at least 24 hours.

The nematodes migrated downward through the tissue paper in the clear water of petri dish. This clear water containing nematodes was taken in a big test tube with the help of a dropper by washing the petri dish several times with clean water, so that no nematodes were left out in the petri dish. The water in test tubes, along with nematodes, was kept undisturbed for 2-3 hours, so that the nematodes in the water could settle down at the bottom of the test tube. Then most of the water in the test tube was withdrawn very carefully without disturbing the nematodes. Finally the entire nematode population along with 3-4 ml of water was taken after decantation of water.

Fixation and Preservation of Nematodes:

The nematodes, kept in 3-4 ml of water in the test tube, were killed and fixed instantly in their characteristic body posture by Seinhorst's method

(Seinhorst, 1966) in hot FA (formalin-acetic acid 4:1) solution, the composition of which was as follows:

Formalin (40% formaldehyde)	: 08 ml
Glacial acetic acid	: 02 ml
Distilled water	: 90 ml

These fixed nematodes were preserved in the same solution in properly labeled specimen tubes.

RESULT

List 1: Systemic Index of Dorylaimid Nematodes Available from Himachal Pradesh

Phylum NEMATODA Rudolphi, 1808
(Lankester, 1877)

Order DORYLAIMIDA Pearse, 1942

Suborder DORYLAIMINA Pearse, 1936

Superfamily DORYLAIMOIDEA De Man, 1976

1. Family DORYLAIMIDAE De Man, 1976

Subfamily DORYLAIMINAE De Man, 1976

1. Genus *Dorylaimus* Dujardin, 1845

1. *Dorylaimus innovatus* Jana and Baqri, 1982

2. *D. neominimus* Gantait *et al.*, 2010

3. *D. thornei* Andrassy, 1969

Subfamily LAIMYDORINAE Andrassy, 1969

2. Genus *Laimydorus* Siddiqi, 1969

4. *Laimydorus* sp.

2. Family APORCELAIMIDAE Heyns, 1965

Subfamily APORCELAIMINAE Heyns, 1965

3. Genus *Aporcelaimellus* Heyns, 1965

5. *Aporcelaimellus* sp.

6. A. *conicaudatus* (Altherr, 1953) Monterio, 1970

7. A. *heyngsi* Baqri and Jairajpuri, 1968

3. Family QUDSIANEMATIDAE

Jauirajpuri, 1965

Subfamily QUDSIANEMATINAE

Jauirajpuri, 1965

4. Genus *Indokochinema* Darekar and

Khan, 1979

8. Genus *Indokochinema* sp.
5. Genus *Labronema* Thorne, 1939
9. *L. glandosum* Rahman *et al.*, 1986
6. Genus *Thonus* Thorne, 1974
10. *T. garhwaliensis* Ahmad *et al.*, 1986
4. Family NORDIIDAE Jairajpuri and Siddiqi, 1964
Subfamily PUNGENTINAE Siddiqi, 1969
7. Genus *Pungentus* Thorne & Swanger, 1936
11. *Pungentus clavatus* Ahmad & Jairajpuri, 1979
8. Genus *Enchodelus* Thorne, 1939
Subgenus *Enchodelus* Thorne, 1939
12. *Enchodelus (Enchodelus) distinctus* Ahmad & Jairajpuri, 1980
13. *E. (E.) microdoroides* Baqri & Jairajpuri, 1974
Subgenus *Nepalus* Ahmad & Jairajpuri, 1980
14. *Enchodelus (Nepalus) maximus* Baqri & Jairajpuri, 1974
Subgenus *Paraenchodelus* Ahmad & Jairajpuri, 1980
15. *Enchodelus (Paraenchodelus) longidens* Jairajpuri & Loof, 1968
16. *E. (P.) satendri* Baqri & Jairajpuri, 1968
17. *E. (P.) thornei* Baqri & Jairajpuri, 1974
Subgenus *Rotundus* Ahmad & Jairajpuri, 1980
18. *Enchodelus (Rotundus) parateres* Baqri & Jairajpuri, 1974
Subfamily ACTINOLAIMOIDINAE Meyl, 1957
9. Genus *Actinolaimoides* Meyl, 1957
19. Genus *Actinolaimoides* sp.
Superfamily LONGIDOROIDEA Thorne, 1935
5. Family LONGIDORIDAE Thorne, 1935
Subfamily LONGIDORINAE Thorne, 1935
10. Genus *Paralongidorus* Siddiqi, Hooper & Khan, 1963
20. Species *Paralongidorus citri* (Siddiqi, 1959)
Siddiqi, Hooper & Khan, 1963
Syn. *Xiphinema citri* Siddiqi, 1959
11. Genus *Longidorus* Micoletzky, 1922
21. Species *Longidorus himalayensis* (Khan, 1986) Jairajpuri & Ahmad, 1992
Syn. *Neolongidorus himalayensis* Khan, 1986
22. *Longidorus elongates* (De Man, 1876)
Thorne & Swanger, 1936
Dorylaimus elongatus De Man, 1876
6. Family XIPHINEMATIDAE Dalmasso, 1969
Subfamily XIPHINEMATINAE Dalmasso, 1969
12. Genus *Xiphinema* Cobb, 1913
23. *Xiphinema* sp.
24. *Xiphinema americanum* Cobb, 1913
25. *X. basiri* Siddiqi, 1959
Syn. *X. cobbi* Sharma & Saxena, 1981
Syn. *X. hayati* Javed, 1983
26. *X. bergeri* Luc, 1973
Superfamily BELONDIROIDEA Thorne, 1939
7. Family BELONDIRIDAE Thorne, 1939
Subfamily BELONDIRINAE Thorne, 1939
- 7.13. Genus *Axonchium* Cobb, 1920
Subgenus *Axonchium* Cobb, 1920
27. *Axonchium (Axonchium) manalicum* Ali, Jairajpuri & Coomans, 1975
Subfamily DORYLAIMELLINAE Jairajpuri, 1964
14. Genus *Dorylaimellus* Cobb, 1913
Subgenus *Dorylaimellus* Cobb, 1913
28. *Dorylaimellus (Dorylaimellus) himalayensis* Ali, Jairajpuri & Coomans, 1974
Superfamily TYLENCHOLAIMOIDEA Filipjev, 1934
8. Family TYLENCHOLAIMIDAE Filipjev, 1934
Subfamily TYLEPTINAE Jairajpuri, 1964
15. Genus *Gymnotyleptus* Ahmad & Jairajpuri, 1982
29. *Gymnotyleptus gymnochilus* (Loof, 1964)
Ahmad & Jairajpuri, 1982
Syn. *Tyleptus gymnochilus* Loof, 1964

30. *Gymnotyleptus indicus* Ahmad & Jairajpuri, 1982
 Superfamily NYGOLAIMOIDEA Thorne, 1935
9. Family NYGOLAIMIDAE Thorne, 1935
 Subfamily NYGOLAIMINAE Thorne, 1935
16. Genus *Aquatides* Heyns, 1968
31. *Aquatids christicki* Ahmad & Jairajpuri, 1982
32. A. deconincki Jairajpuri & Coomans, 1977
17. Genus *Clavicaudoides* Heyns, 1968
33. *Clavicaudoides tenuicaudatum* Ahmad & Jairajpuri, 1982
18. Genus *Paravulvus* Heyns, 1968
34. Species *Paravulvus papillatus* Ahmad & Jairajpuri, 1982
7. *C. costatus* (De Man, 1921) Siddiqi, 1978
8. *C. indicus* (Khan, Chawla & Prasad, 1969) Siddiqi, 1978
 Syn. *Tylenchus (Aglenchus) indicus* Khan, Chawla & Prasad, 1969
4. Genus *Filenchus* Andrassy, 1954
 (Meyl, 1961)
9. Species *Filenchus* sp.
5. Genus *Polenchus* Andrassy, 1980
10. Species *Polenchus shamimi* Baqri, 1991
 Subfamily BOLEODORINAE Khan, 1964
6. Genus *Boleodorus* Thorne, 1941
11. Species *Boleodorus mirus* Khan, 1964
 Subfamily DUOSULCIINAE Siddiqi, 1979
7. Genus *Malenchus* Andrassy, 1968
 Subgenus *Malenchus* Andrassy, 1968
12. Species *Malenchus (Malenchus)* sp.
8. Genus *Ottolenchus* Husain & Khan, 1967
 (Golden, 1971)
13. Species *Ottolenchus eqisetus* (Husain & Khan, 1967) Wu, 1970
 Syn. *Ottolenchus purvus* (Siddiqi) Siddiqi, 1979
 Subfamily PLEUROTYLENCHINAE Andrassy, 1976
9. Genus *Cephalenchus* Goodey, 1962
 (Geraert, 1968)
14. Species *Cephalenchus leptus* Siddiqi, 1963
 Infraorder ANGUINATA Siddiqi, 2000
- Superfamily ANGUINOIDEA Nicoll, 1935 (1926)
2. Family ANGUINIDAE Nicoll, 1935 (1926)
 Subfamily ANGUININAE Nicoll, 1935 (1926)
10. Genus *Ditylenchus* Filipjev, 1936
15. Species *Ditylenchus myceliophagus* Goodey, 1958
 11. Genus *Nothotylenchus* Thorne, 1941
16. Species *Nothotylenchus* sp.

List 2: Systemic Index of Tylenchid Nematodes Available From Himachal Pradesh

- Class SECERNENTEA von Linstow, 1905
- Subclass TYLENCHIA Inglis, 1983
- Order TYLENCHIDA Thorne, 1949
- A] Suborder TYLENCHINA Chitwood in Chitwood Chitwood, 1950
- Infraorder TYLENCHATA Siddiqi, 2000
- Superfamily TYLENCHOIDEA Örley, 1880
1. Family TYLENCHIDAE Örley, 1880
- Subfamily TYLENCHINAE Örley, 1880
1. Genus *Tylenchus* Bastain, 1865
1. Species *Tylenchus arcuatus* Siddiqi, 1963
2. *T. elegans* De Man, 1876
3. *T. magnus* Khurma & Gupta, 1988
2. Genus *Aglenchus* Andrassy, 1954
 (Meyl, 1961)
4. Species *Aglenchus agricola* (De Man, 1884)
 Meyl, 1961
5. *A. muktii* Phukan & Sanwal, 1980
3. Genus *Coslenchus* Siddiqi, 1978
6. Species *Coslenchus capsici* Khurma & Gupta, 1988

12. Genus *Safianema* Siddiqi, 1980
17. Species *Safianema* sp.
- B] Suborder Hoplolaimina Chizhov & Berezina, 1988
- Superfamily HOPLOLAOMOIDEA Filipjev, 1934 (Paramonov, 1967)
3. Family HOPLOLAIMIDAE Filipjev, 1934
- Subfamily HOPLOLAIMINAE Filipjev, 1934
13. Genus *Hoplolaimus* von Daday, 1905
- Subgenus *Basirolaimus* Shamsi, 1979
18. Species *Hoplolaimus (Basirolaimus) chambus* Jairajpuri & Baqri, 1973
- Syn. *Hoplolaimus chambus* Jairajpuri & Baqri, 1973
- Syn. *Basirolaimus chambus* (Jairajpuri & Baqri, 1973) Siddiqi, 1986
19. *Hoplolaimus (Basirolaimus) indicus*
- Syn. *Hoplolaimus indicus* Sher, 1963
20. *Basirolaimus indicus* (Sher, 1963)
- Shamsi, 1979
14. Genus *Scutellonema* Andrassy, 1958
21. Species *Scutellonema brachyurus* (Steiner, 1938) Andrassy, 1958
- Syn. *Rotylenchus brachyurus* Steiner, 1938
- Subfamily ROTYLENCHOIDINAE Whitehead, 1958
15. Genus *Rotylenchoides* Whitehead, 1958
22. *Rotylenchoides neoformis* (Siddiqi and Husain, 1964) Sher, 1966
16. Genus *Helicotylenchus* Steiner, 1945
23. Species *Helicotylenchus borinquensis* Román, 1965 (*sp. inq.* For Fortuner *et al.*, 1981)
24. *H. girus* Saha *et al.*, 1973
25. *H. goldeni* Sultan & Jairajpuri, 1979
26. *H. rohtangus* Jairajpuri & Baqri, 1973
- (Syn. of *H. retusus* for Lal & Khan, 1997)
27. *H. rotundicaudata* Sher, 1966 (Syn. Of *H. dihystera* for Fortuner *et al.*, 1981)
28. *H. shakili* Sultan, 1981
29. *H. vulgaris* Yuen, 1964
17. Genus *Orientylus* Jairajpuri & Siddiqi, 1977
30. Species *Orientylus geraerti* Jairajpuri & Siddiqi, 1979
- Syn. *Rotylenchus geraerti* (Jairajpuri & Siddiqi) Zancada & Lima, 1986
18. Genus *Rotylenchus* Filipjev, 1936
- Subgenus *Rotylenchus* Filipjev, 1936
31. Species *Rotylenchus (Rotylenchus) dalhousiensis* Sultan & Jairajpuri, 1979
32. *R. (R.) indorobustus* Jairajpuri & Baqri, 1973
- (Syn. of *R. robustus* for Brzeski, 1998)
33. *R. (R.) neorobustus* Sultan & Jairajpuri, 1979
34. *Rotylenchus (Rotylenchus)* sp.
4. Family ROTYLENCHULIDAE Husain and Khan, 1957 (Husain, 1976)
- Subfamily ROTYLENCHULINAE Husain and Khan, 1967
19. Genus *Rotylenchulus* Linford and Oleiveira, 1940
35. Species *Rotylenchulus reniformis* Linford and Oleiveira, 1940
5. Family PRATYLENCHIDAE Thorne, 1949 (Siddiqi, 1963)
- SubFamily PRATYLENCHINAE Thorne, 1949
20. Genus *Pratylenchus* Filipjev, 1936
36. Species *Pratylenchus coffeae* (Zimmermann, 1898) Filipjev & Schurr. Stekh., 1941
- Syn. *Tylenchus coffeae* Zimmermann, 1898
37. *P. neglectus* (Rensch, 1924) Filipjev & Sch. Stekh., 1941
- Syn. *Aphelenchus neglectus* Rensch, 1924
- Syn. *Tylenchus neglectus* (Rensch) Steiner, 1928.
38. *P. neocapitatus* Khan & Singh, 1975
39. *P. penetrans* (Cobb, 1917) Filipjev & Sch. Stekh., 1941
- Syn. *Tylenchus penetrans* Cobb, 1917

40. *P. zae* Graham, 1951
 Syn. *P. impar* Khan & Singh, 1975
41. *Pratylenchus* sp.
 Subfamily HIRSCHMANNIELLANAE
 Fotedar & Handoo, 1978
21. Genus ***Hirschmanniella*** Luc &
 Goodey, 1964
42. Species *Hirschmanniella mucronata* (Das,
 1960) Khan, Siddiqi, Khan, Husain &
 Saxena, 1964
 Syn. *Radhopholus mucronata* Das, 1960
43. *H. oryzae* (Van Breda de Hann, 1902) Luc
 & Goodey, 1964
 Syn. *Tylenchus oryzae* Van Breda de Hann,
 1902
6. Family MELOIDOGYNIDAE Skarbilovich,
 1959 (Wouts, 1973)
 Subfamily MELOIDOGYNIDAE
 Skarbilovich, 1959
22. Genus ***Meloidogyne*** Goeldi, 1892
44. Species *Meloidogyne arenaria* (Neal, 1889)
 Chitwood, 1949
 Syn. *Anguina arenaria* Neal, 1889
45. *M. graminicola* Goloden & Birchfield, 1965
46. *M. hapla* Chitwood, 1949
47. *M. incognita* (Kofoid & White, 1919)
 Chitwood, 1949
 Syn. *Oxyuris incognita* Kofoid & White, 1919
 Syn. *Heterodera incognita* (Kofoid & White)
 Sandground, 1923
48. *M. javanica* (Treub, 1885) Chitwood, 1949
 Syn. *Heterodera javanica* Treub, 1885
7. Family HETERODERIDAE Filipjev &
 Schurr. Stekh., 1941 (Skarbilovich, 1947)
 Subfamily HETERODERINAE Filipjev &
 Schurr. Stekh., 1941
23. Genus ***Heterodera*** Schmidt, 1871
49. Species *Heteroder aavenae* Wollenweber, 1924
50. *H. trifolii* Goffart, 1932
 24. Genus ***Globodera*** Skarbilovich, 1959
51. Species *Globodera pallida* (Stone, 1973)
 Behrens, 1975
 Syn. *Heterodera pallida* Stone, 1973
52. *G. pseudorostochiensis* (Kirjanova, 1963)
 Mulvey & Stone, 1976
 Syn. *G. rostochiensis* Krall, 1978
 Syn. *Heterodera pseudorostochiensis*
 Kirjanova, 1963
 Superfamily DOLICHODOROIDEA Chitwood
 in Chitwood & Chitwood, 1950 (Siddiqi, 1968)
8. Family TELOTYLENCHIDAE Siddiqi, 1960
 Subfamily TELOTYLENCHINAE Siddiqi, 1960
25. Genus ***Quinisulcius*** Siddiqi, 1971
53. Species *Quinisulcius capitatus* (Allen, 1955)
 Siddiqi, 1971
 Syn. *Tylenchorhynchus capitatus* Allen, 1955
54. *Q. indicus* Luqman & Khan, 1986
26. Genus ***Tylenchorhynchus*** Cobb, 1913
55. Species *Tylenchorhynchus clarus* Allen, 1955
56. *T. cylindricus* Cobb, 1913
57. *T. mashhoodi* Siddiqi & Basir, 1959
 Syn. *Macrorhynchus mashhoodi* (Siddiqi &
 Basir) Sultan, Singh & sakuja, 1991
58. *Tylenchorhynchus* sp.
27. Genus ***Bitylenchus*** Filipjev, 1934
 Syn. *Tylenchus* (Bitylenchus Filipjev, 1934)
59. Species *Bitylenchus brevilineatus* (Williams,
 1960) Jairajpuri, 1982
 Syn. *Tylenchorhynchus brevilineatus*
 Williams, 1960
60. *Bitylenchus* n. sp.
 Subfamily MERLINIINAE Siddiqi, 1971
28. Genus ***Merlinius*** Siddiqi, 1970
61. Species *Merlinius nothus* (Allen, 1955)
 Siddiqi, 1970
 Syn. *Tylenchorhynchus nothus* Allen., 1955

62. *M. nizami* Luqman & Khan, 1986
63. *Merlinius* sp.
9. Family PSILENCHIDAE Paramonov, 1967
(Khan, 1969)
- Subfamily PSILENCHINAE Paramonov, 1967
29. Genus *Psilenchus* De Man, 1921
64. Species *Psilenchus aestuarius* Andrassy, 1962
65. *P. hilarulus* De Man, 1921
66. *P. hilarus* Siddiqi, 1963
- Syn. *Psilenchus neoformis* Jairajpuri & A. H. Siddiqi, 1963
- C] Suborder CRICONEMATINA Siddiqi, 1980
- Superfamily CRICONEMATOIDEA Taylor, 1936 (1914) Geraert, 1966
10. Family CRICONEMATIDAE Taylor, 1936 (1914) Thorne, 1949
- Subfamily CRICONEMATINAE Taylor, 1936 (1914)
30. Genus *Ogma* Southern, 1914
- Subgenus *Ogma* Southern, 1914
67. Species *Ogma (Ogma) simlaensis* (Jairajpuri, 1963) Andrassy, 1979
- Syn. *Variasquamata rhosimum* Khan, Chawla & Saha., 1976
31. Genus *Macroposthonia* De Man, 1880
68. Species *Macroposthonia bilaspurensis* Gupta & Gupta, 1981
69. *M. curvata* (Raski, 1952) De Grisse & Loof, 1965
- Syn. *Criconemella curvata* Raski, 1952
70. *M. xenoplax* (Raski, 1952) De Grisse & Loof, 1965
- Syn. *Criconemoides xenoplax* Raski., 1952
- Subfamily HEMICRICONEMOIDINAE Andrassy, 1979
32. Genus *Hemicriconemoides* Chitwood & Birchfield, 1957
71. Species *Hemicriconemoides gaddi* (Loos, 1949) Chitwood & Birchfield, 1957
- Syn. *Criconemoides gaddi* Loos., 1949
72. *H. mangiferae* Siddiqi, 1961
- Syn. *H. litchi* Edward & Misra., 1964
- Syn. *H. aberrans* Phukan & Sanwal., 1983
- Superfamily HEMICYCLIOPHOROIDEA Skarbilovich, 1959 (Siddiqi, 1980)
11. Family HEMICYCLIOPHOROIDAE Skarbilovich, 1959 (Geraert, 1966)
- Subfamily HEMICYCLIOPHOROINAE Skarbilovich, 1959
33. Genus *Hemicycliophora* De Man, 1921
73. Species *Hemicycliophora subaolica* Jairajpuri & Baqri, 1973
74. *Hemicycliophora* sp.
- Superfamily TYLENCHULOIDEA Skarbilovich, 1947 (Raski & Siddiqi, 1975)
12. Family TYLENCHULOIDAE Skarbilovich, 1947 (Kirjanova, 1955)
- Subfamily TYLENCHULOINAE Skarbilovich, 1947
34. Genus *Tylenchulus* Cobb, 1913
75. Species *Tylenchulus semipenetrans* Cobb, 1913
13. Family PARATYLENCHIDAE Thorne, 1949 (Raski, 1962)
- Subfamily PARATYLENCHINAE Thorne, 1949
35. Genus *Paratylenchus* Micoletzky, 1922
- Subgenus *Gracilacus* Raski, 1962 (Siddiqi, 1986)
- Syn. *Gracilpaurus* Ganguly & Khan., 1990
76. Species *Paratylenchus (Gracilacus) aonli* Misra & Edward, 1971
- Syn. *Gracilacus aonli* (Misra & Edward) Raski., 1976
- Subgenus *Paratylenchus* Micoletzky, 1922
77. *Paratylenchus (Paratylenchus) colbrani* *colbrani* Raski, 1975

78. *P. (P.) elachistus* Steiner, 1949
79. *P. (P.) lepidus* Raski, 1975
80. *P. (P.) minor* Sharma & Khan, 1986
81. *P. (P.) nanus* Cobb, 1923
82. *P. (P.) runii* Sharma & Khan, 1986
83. *P. (P.) vandenbrandei* De Grisse, 1962

DISCUSSION

The study on nematodes started in Himachal Pradesh with the report of Thirumalaha (1951) about the infestation of root-knot nematode on potato tubers for the first time from Shimla and subsequently, the first nematology unit was established at Central Potato Research Institute, Shimla. A total of 138 nematode species under 63 genera belonging to Dorylaimida, Tylenchida and other nematode orders have been reported from Himachal Pradesh of which 45 species were described as new (Sharma & Gupta, 1998). Sharma & Kaur (1987) observed the parasitism of root-knot nematode *Meloidogyne* sp. on apple which is one of the most economically important cash crops of Himachal Pradesh. The species of root-knot nematodes like *Meloidogyne incognita*, *M. hapla* and *M. javanica* are the most prominent pest of potato and this nematode has been found to be prevalent in about 60% of the potato growing areas of Himachal Pradesh (Krishna Prasad, 1986) and prevalence and intensity of root-knot nematode infection on potato has also been observed (Krishna Prasad & Rajendran, 1986). *Quinisulcius capitatus* has also been suspected as potential pest of potato (Krishna Prasad, 1986; Khan et al., 1990). The infestation of *Meloidogyne* spp. on tomato in Solan area has also been observed (Bharadwaj et al., 1974). *Meloidogyne incognita* and *M. hapla* are the only recognized pests of vegetable crops in the state and these nematodes have been reported to occur in 50 – 100% of vegetable fields of Himachal Pradesh (Bharadwaj et al., 1974; Reddy, 1989). *Tylenchorhynchus* and *Helicotylenchus* have been reported to occur on ginger being highly prevalent (Kaur et al., 1989). Saharan and Chohan (1972) surveyed the disease of plants due to nematodes in Kangra district of

the state. *Tylenchorhynchus* spp., *Macroposthonia xenoplex*, *Meloidogyne incognita*, *Pratylenchus pratensis*, *Helicotylenchus* spp., *Paratylenchus* sp. and *Xiphinema* sp. have been observed to occur on agricultural, horticultural and forest trees (Sharma et al., 1982, Anonymous, 1993; Chandel, 1993). *Meloidogyne incognita*, *M. hapla*, *Pratylenchus penetrans*, *P. coffeae*, *Paratylenchus prunii*, *Tylenchulus semipenetrans*, *Macroposthonia xenoplex* and *Ditylenchus myceliophagus* are the most important nematode species which cause enormous yield losses in various economically important agricultural and horticultural crops and the wide spread occurrence of *Heterodera zeae*, *Quinisulcius capitatus*, *Tylenchorhynchus*, *Helicotylenchus* and *Xiphinema* in a number of agricultural and horticultural crops also cause direct or indirect crop losses (Sharma & Gupta, 1998). In the present study, 34 species under 18 genera and 9 families of Dorylaimida and 83 species under 35 genera and 12 families of Tylenchida have been compiled and reported. Among these, *Dorylaimus neominimus* Gantait et al. 2010, *Laimydorus* sp., *Aporcelaimellus heynsi* Baqri and Jairajpuri, 1968., *Indokochinema* sp., *Labronema glandosum* Rahman et al., 1986, *Thonus garhwaliensis* Ahmad et al., 1986, *Actinolaimoides* sp. are new rerecords from Himachal Pradesh and *Dorylaimus thornei* Andrassy, 1969 is the first report from India. Five species of Tylenchida i.e. *Polenchus shamimi* Baqri, 1991, *Filenchus* sp., *Rotylenchooides neoformis* (Siddiqi and Husain, 1964) Sher, 1966, *Rotylenchulus reniformis* Linford and Oliveira, 1940 and *Tylenchorhynchus cylindricus* Cobb, 1913 are new records from Himachal Pradesh.

SUMMARY

The present study includes two systematic lists of soil free living and plant parasitic nematodes belonging to the orders Dorylaimida Pearse, 1942 and Tylenchida Thorne, 1949 from Himachal Pradesh. 34 species under 18 genera and 9 families of Dorylaimida and 83 species under 35 genera and 12 families of Tylenchida have been compiled and reported. Among these, *Dorylaimus neominimus* Gantait et al. 2010, *Laimydorus* sp.,

Aporcelaimellus heynsi Baqri and Jairajpuri, 1968., *Indokochinema* sp., *Labronema glandosum* Rahman et al., 1986, *Thonus garhwaliensis* Ahmad et al., 1986, *Actinolaimoides* sp. are new rerecords from Himachal Pradesh and *Dorylaimus thornei* Andrassy, 1969 is the first report from India. Five species of Tylenchida i.e. *Polenchus shamimi* Baqri, 1991, *Filenchus* sp., *Rotylenchoides neoformis* (Siddiqi and Husain, 1964) Sher, 1966,

Rotylenchulus reniformis Linford and Oleiveira, 1940 and *Tylenchorhynchus cylindricus* Cobb, 1913 are new records from Himachal Pradesh.

ACKNOWLEDGEMENT

The authors are grateful to Dr. K. Venkataraman, Director, Zoological Survey of India, Kolkata for providing facilities to carry out the work and to publish the result.

REFERENCES

- Ahmad, M and Jairajpuri, M.S. 1979. Nematodes from high altitude in India. VIII. *Pungentus clavatus* n. sp. with a key to species of the genus *Pungentus* (Nematoda: Dorylaimida). *Revue de Nematologie*, **2**(2): 185-189.
- Ahmad, M and Jairajpuri, M.S. 1980. A review of the genus *Enchodelus* Thorne, 1939 with description of species from India. *Records of Zoological Survey of India, Occasional Paper No.*, **15**: 1-42.
- Ahmad, M and Jairajpuri, M.S. 1982. Some studies on Tyleptinae Jairajpuri, 1964 with a proposal of *Gymnotyleptus* n. gen. (Nematoda: Dorylaimida) *Kanpur University Research Journal* (1980), **1**: 70-76.
- Ali, S.S., Jairajpuri, M.S. and Coomans, A. 1974. Nematodes of high altitude in India. IV. Two new species of Belondiridae Thorne, 1939 (Dorylaimida). *Nematologica*, **19**(1973): 498-507.
- Anonymous, 1993. Annual Report. Reg. Hort. Res. Sta. (UHF) Jachh (Nurpur), Kangra (HP).
- Baqri, Q.H. and Jairajpuri, M.S. 1974. Nematodes of high altitudes in India. V. Five new species of the genus *Enchodelus* Thorne, 1939 (Dorylaimida). *Nematologica*, **20**(2): 131-136.
- Baqri, Q.H. and Jairajpuri, M.S. 1975. Nematodes of high altitudes in India. VI. *Enchodelus thornei* sp. n. and relationship of the genera *Enchodelus*, *Oonaguntus* and *Malekus*. *Indian J. Nematol.*, **5**(2): 185-188.
- Bhardwaj, A.K. and Sharma, P.L. 1971. Preliminary studies on nematodes associated with fruit plants in H.P. *Himachal J. Agric. Res.*, **1**: 53-54.
- Bhardwaj, A.K., Sharma, P.L. and Phunchog, S. 1974. Root knot nematodes (*Meloidogyne* sp.) infestation on tomato in Solan area of Himachal Pradesh. *Indian J. Nematol.*, (1972), **2**: 197-198.
- Bhardwaj, A.K., Sharma, P.L. and Thakur, J.R. 1973. Mushroom nematodes. *Curr. Sci.*, **42**(7): 257-258.
- Chandel, Y.S. 1982. Studies on nematodes associated with mushroom and their control. *M.Sc. Thesis*, HPKV, Solan, H.P.
- Chandel, Y.S. 1986. Studies on the nematodes associated with citrus in Himachal Pradesh. *Ph.D. Thesis*, UHF, Solan (H.P.).
- Chandel, S.S. 1993. Studies on the nematodes associated with *Gladiolus hortulances* L. *M.Sc. Thesis*, UHF, Solan (H.P.)
- Christie, J.R. and Perry, V.G. 1951. Removing nematodes from soil. *Proceedings of Helminthological Society of Washington*, **17**: 106-108.

- Cobb, N.A. 1918. Estimating the nema population of soil. *Agric. Tech. Cir. US Dept. Agric.*, **1**: 48pp.
- Dalal, M.R. and Bhatti, D.S. 1983. Dhaincha (*Sesbania aclecula*) an additional host of *Heterodera cajini* Koshy, 1967. *Proc. 3rd. Nematol. Symp.*, Himachal Pradesh Krishi Viswa Vidlaya, Solan (India). May 26-29, 1983: 10.
- Jairajpuri, M.S. and Ahmad, W. 1992. *Dorylaimida. Freeliving, Predaceous and Plant parasitic Nematodes*. Oxford & IBH Publishing Co. Pvt. Ltd., New Delhi. 458pp.
- Jairajpuri, M.S. and Baqri, Q.H. 1973. Nematodes of high altitude in India. I. Four new species of Tylenchida. *Nematologica*, **19**: 19-30.
- Jairajpuri, M.S. and Coomans, A. 1977. Nematodes from high altitude of India. VII. *Aquatides deconicki* sp. n. (Dorylaimida: Nematoda). *Indian J. Nematol.* (1975), *Indian J. Nematol.*, **5**: 243-246.
- Jairajpuri, M.S. and Siddiqi, A.H. 1963a. On *Psilenchus neoformis* n. sp. (Nematoda: Tylenchida) from Solan (H. P.), North India. *Curr. Sci.*, **32**: 318-319.
- Jairajpuri, M.S. and Siddiqi, A.H. 1963b. *Xiphinema brevicole* Lordello & Da costa, 1961 (Nematoda: Dorylaimoidea) from Dalhousie (H. P.), North India. *Curr. Sci.*, **32**: 508.
- Jairajpuri, M.S. and Siddiqi, M.R. 1979. Observation on the nematode genera *Orientylus* and *Calvitylus* (Rotylenchoidinae: Hoplolaimidae) with description of three new species. *Indian J. Nematol.*, **7**: 101-111.
- Kaur, D.J. 1987. Studies on nematodes associated with ginger (*Zingiber officinale* Rosc.) in Himachal Pradesh. *Ph.D. Thesis*, HPU, Shimla (H.P.).
- Kaur, D.J. and Sharma, N.K. 1990. A new report of *Pratylenchus coffeae* – A cause of ginger yellows. *Int. Nematol. Network Newslet.* **7**(1): 15-16.
- Kaur D.J., Sharma, N.K. and Kalia, D.C. 1989. Nematodes associated with ginger (*Zingiber officinale* Rosc.) in Himachal Pradesh. *Indian J. Hill Farmg.* **3**(1): 49-51.
- Kaur D.J., Sharma, N.K. and Khan, M.L. 1990. Occurrence of *Pratylenchus zeae* Graham, 1951 on ginger (*Zingiber officinale* Rosc.) in Himachal Pradesh. *Indian J. Nematol.*, **19**: 68.
- Khan, E. 1964. *Boleodorus mirus* n. sp. (Tylenchida: Boleodorinae n. subfam.) from Kufri, Simla (H. P.), India with a key to species of the genus *Boleodorus* Thorne, 1941 *Zool. Anz.*, **179**: 336-341.
- Khan, E., Ganguly, S. and Lalitha, Y. 1990. *Quinisulcius capitatus* - a potential pest of potato in Himachal Pradesh, India. *Int. Nematol. Network Newslet.* **7**(4): 22-24.
- Khan, E. and Singh, D.B. 1974. Five new species of *Pratylenchus* (Nematoda: Pratylenchidae) from India. *Indian J. Nematol.*, **4**(2): 199-211.
- Khan, M.L. and Khan, S.H. 1982. Three new species of the genus *Orientylus* Jairajpuri and Siddiqi, 1977 associated with fruit trees in India (Rotylenchoidinae: Nematoda). *Indian J. Nematol.*, **12**(1): 111-117.
- Khurma, U.R. 1989. Plant parasitic nematodes from some hilly areas of North India. *Indian J. Nematol.*, **18**: 132-133.
- Khurma, U.R. Gupta, N.K. 1988a. Description of *Tylenchus magnus* n. sp. (Nemata: Tylenchidae). *Revue de Nematologie*, **11**(1): 35-37.
- Khurma, U.R. Gupta, N.K. 1988b. *Coslenchus capsici* n. sp. (Nematoda: Tylenchidae) from India. *Systematic Parasitology*, **11**(3): 227-229.

- Krishna Prasad, S.K. 1986. Problens of potato. In: *Plants Parasitic Nematodes of India – Problems and Progress* (Edited by G. Swarup and D.R. Dasgupta). IARI, New Delhi, India : 350-370.
- Krishna Prasad, S.K. and Rajendran, T.P. 1986. Prevalence and intensity of root-knot nematode infection on potato in Himachal Pradesh. *Indian Int. Nematol. Network Newslet.*, **3**(3): 13-15.
- Mukhopadhyaya, M.C. 1970. Nematodes associated with fruit trees and vegetable crops in Himachal Pradesh. *J. Res.*, PAU, **7**: 656-658.
- Reddy, M.C.M. 1989. Fungi as biological control agents of root-knot nematodes. *Ph.D. Thesis*, UHF, Solan (H.P.).
- Saha, M, Chawla, M.L. and Khan, E. 1973. Two new species of *Helicotylenchus* Steiner, 1947 from North India. *Indian J. Nematol.*, **3**(2): 83-87.
- Saharan, G.S. and Chohan, J.S. 1972. A survey of disease of plants in Kangra district, Himachal Pradesh. *J. Res.*, **9**: 68-73.
- Seinhorst, J.W. 1966. Killing nematodes for taxonomic study with hot f. a. 4:1. *Nematologica*, **1**: 178.
- Sharma, G.C., Khan, E. and Sharma, N.K. 1986. Two new species of the genus *Paratylenchus* Micoletzky, 1922 (Nematoda: Paratylenchidae) from Himachal Pradesh, India. *Indian J. Nematol.*, **16**: 231-235.
- Sharma, G.S., Sharna, N.K. and Khan, M.L. 1988. Plant parasitic nematodes of plum (*Prunus domestica*) in Himachal Pradesh. *Indian J. Hort.* **45**: 355-358.
- Sharma, N.K., Damaneet and Sharma, S.B. 1984. Incidence of maize cyst nematodes, *Heterodera zeae* in Himachal Pradesh. *Indian J. Nematol.*, **14**: 198.
- Sharma, N.K. and Gupta, R. 1998. Plant nematology in Himachal Pradesh: An overview. In: *Phytonematology in India* (edited by Pravin Chandra Trivedi), CBS Publishers and Distributors, New Delhi, India : 33-46.
- Sharma, N.K. and Kaur, D.J. 1986. Nematodes associated with temperate fruit crops in Himachal Pradesh. In: *Advances in Research on temperate fruit* (edited by Chadha, T.R., Bhutani, V.P., Kaul, J.L.), Solan, Himachal Pradesh, India, Dr. Y.S. Parmar Univ. of Horticulture and Forestry: 359-362.
- Sharma, N.K. and Kaur, D.J. 1987. Parasitism of root-knot nematode. *Meloidogyne* sp. on apple in Himachal Pradesh (India). *Indian J. Hill. Farmg.* **1**: 69.
- Sharma, N.K., Nath, A. and Thapa, C.D. 1982. Nematodes associated with forest nurseries. *Proc. Improvement in Forest Biomass*, UHF, Solan (HP) : 459-461.
- Siddiqi, M.R. 2000. *Tylenchida, parasites of plants and insects*, 2nd. edition. CABI Publishing, CAB International, Wallingford, U.K. xvii + 833pp.
- Sultan, M.S. 1980. Two new species of the genus *Orientylus* Jairajpuri and Siddiqi, 1977 (Tylenchida: Rotylenchoidinae). *Revue de Nematologie*, **3**(2): 227-229.
- Sultan, M.S. and Jairajpuri, M.S. 1978. Nematodes of high altitude in India. III. *Helicotylenchus goldeni* n. sp. (Nematoda: Tylenchida). *Indian J. Nematol.*, **8**(2): 169-171.
- Sultan, M.S. and Jairajpuri, M.S. 1979. Nematodes of high altitude in India. IX. Description of two new species of *Rotylenchus* (Nematoda: Tylenchida). *Revue de Nematologie*, **2**(2): 191-196.
- Thirumalachar, M.J. 1951. Root-knot nematode on potato tubers in Shimla. *Curr. Sci.*, **20**(4): 104.